Real-time ground motion simulation and visualisation

Current readme for real-time ground motion simulation

1. Source generation (Viktor/Hoby)

- a. Srf Generation/Plotting
- Manually extract location, dip, strike, rake from Mw tensor (Ristau) into setSrfParams.py
- Change TYPE=2 (point source to finite fault)
- Run createSRF.py

2. Velocity model generation

- a. Preliminary steps to setup on hypocentre (if required)
 - i. Clone the AutoVMGen repo https://github.com/ucgmsim/Auto-Vel-Mod-Generation.git
 - ii. Clone the VM repo into the AutoVMGen directory (or create a symlink) https://github.com/ucgmsim/Velocity-Model.git
 - 1. Compile the VM code by executing 'make parallel'
- b. Run generate_parameters.py
 - i. Uses hypocentre Mw, depth, lat, lon (and others optional inputs e.g. hh)
 - ii. creates a file called params_vel.py which has the velocity model parameters (Lx, Ly, Lz, hh, model origin, model rotation) and simulation duration
- c. Run investigateDomain.py
 - i. Loads the parameters stored in params_vel.py
 - ii. Outputs a plot of the VM domain on a map (found in the directory 'Domain')
 - iii. A simulation duration estimate is printed to the screen (assuming 512 core simulation)
 - iv. If a change in velocity model parameters is required, update desired variables in params_vel.py, run validateparameters.py and then rerun investigateDomain.py (can repeat this step until satisfied with the VM domain)
- d. Run genDomain.py
 - i. Generates the VM and stores it the directory Rapid_Domain, with extracted slices and domain on map plots.

1. Observed ground motions

a. Ahsan

NOTE: Installation is required only once, unless there is a need to update. geoNet package is self contained and should work on your pcs and laptops too.

0) INSTALL geoNet. From https://github.com/ucgmsim/Pre-processing/tree/master/geoNet clone geoNet and follow the installation instructions.

 b. Copy event_info.txt found at geoNet/examples and modify. Then do the following: from geoNet import realtime realtime.run("event_info.txt")

Alternatively performs the steps (1) to (5) outlined below.

1) GET THE REQUIRED SCRIPTS. From geoNet/examples copy

getData.py event_statsll.py processData.py plot_accvel.py plot_psa.py

to your working directory

2) GET THE DATA FOR THE EARTHQUAKE.

In getData.py set BASE_URL to the location to the geoNet URL e.g BASE_URL="ftp://ftp.geonet.org.nz/strong/processed/Proc/2017/02_Feb/2017-02-01_102129/Vol1/data/"

save and run.

 GET THE LIST OF STATIONS FOR THE EARTHQUAKE Run event_statsII.py which creates event_stats.II and places it in your working directory.

4) PROCESS THE GEONET VOL1 DATA TO GET NORTH, EAST AND VERTICAL COMPONENTS Run processData.py which filters and rotates the SMS data placing them in Vol1/data/accBB etc. Note: It is better to redirect the std output to a file like so python processData.py > processData_output

5) CREATE PLOTS FOR OBSEREVED DATA

- Run plot_accvel.py and plot_psa.py
- 2. Ground motion simulation computation (Sung)
 - a. Upload Rupture model to /nesi/projects/nesi00213/RupModel/Realtime/MonYEAR where MonYear is like Dec2016, Jan2017. Under this directory, you need Srf and Stoch.
 - b. Upload Velocity model to /nesi/projects/nesi00213/VelocityModels/. Currently we have Canterbury or South_Island. Unless it is a Canterbury specific model, place it under South_Island. We will be doing some clean up later. Your velocity model should include rho3dfile.d, vp3dfile.p, vs3dfile.s, params_vel.py and outputs from gen_cords (gridfile, gridout, model_bounds, model_coords, model_params).
 - c. Upload station files to /nesi/projects/nesi00213/StationInfo. This should include geonet_stations_xxx.ll, .vs30 and .vs30ref.
 - d. Go to /nesi/projects/nesi00213/RunFolder and execute ./install.sh. and follow the instruction.
 - e. During install.sh, it will advise you expected wallclock time. Enter a sensible wall clock time.
 - f. Follow the remaining instruction.

3. simulation post-processing

- a. Viktor/Hoby
 - b. General Plotting Instructions: run on hypocentre, not fitzroy make sure the scripts are in PATH (hypocentre): export PATH=\$PATH:/nesi/projects/nesi00213/gm_sim_workflow/devel c. Plotting Station based data such as IMs, Vs30, Observed PGA, pSA:
 - plot_stations.py datafile.ll Run on hypocentre, datafile has 6 line header as described at the top of plot stations.py

- d. Plotting Timeslice based data (includes PGV, MMI, timeslice animation) on hypocentre: plot_transfer.py auto /nesi/projects/nesi00213/RunFolder/baes/RT2017Jan17_VMSI_20170117_drill_200mh0p200 EMODv3p0p4 170116. If you have a different username on fitzroy then you will need to add a 4th parameter, your username on fitzroy will copy the remote fitzroy folder to the current directory (.) and start plotting using defaults run plot_ts_sum.py to re-run PGV and MMI plots
 - run plot_ts.py to re-run timeslice image creation

Key lessons from past drills:

1. March 2017 drill (2014 Mw6.3 Eketahuna Earthquake)

Task 1. Source (HOBY)	Expected time reqd.	P e o ple	Notes
GeoNet Mw email to SRF (a) from Mw tensor get Rup parameters (b) From Rup parameters use createSRF.py to make rupture SRF and to plot (c) Execution of createSRF.py	1minute 25sec 2min total	H o b y ∕V ik tor	 a) Manually extract location, dip,strike, rake from Mw tensor (Ristau) into setSrfParams.py b) Change TYPE=2 (point source to finite fault) Use either version "3.3" or "5.3.2a" c) Run "createSRF.py" Please note that the execution time of createSRF.py may change based on the event size

2. Velocity Model Domain (ETHA	N/ROBIN)		
From rupture location to VM dev and domain (a) From location and Mw determine VM origin, 3D extent, and simulation duration (b) Plot VM domain extent (c) Generate VM with H=0.1km for this domain, then extract slices	March Drill Setup code - 4 mins generateDo main initial (inc. getting CMT data) - 2+2=4 min creating VM (inc. manual adjustment) - 10+1 = 11 genDomain - 6+6+6 = 18 Total ~37- 40 minutes	Et h a n, R o bi n, 	March Drill (Robin) Issues encountered/comments 1. nx and ny were incorrectly doubled (already fixed). 2. Development oversights leaving out the writing of params, and importing of shared. Overall if everything went smoothly and no manual adjustment was required, this would total something around 15 minutes. (note this is also dependent on the VM domain size/source magnitude).
From non-uniform station grid, get statgrid points (a) Locations of statgrid points (b) Locations of GeoNet strong motion stations (c) From NZ Vs30 model get Vs30 values at all points (a) Plot VM domain extent, stations (statgrid and GeoNet), (b) color based on Vs30 values	(ignore for now) 1-2 hours!! 10mins -2hours 2mins	A h s a n, C hrr is (a) Vi kt or (b) N ot d o ne	[need .II file for all of NZ (Ahsan)] (Done) run statcords (to get station locations corresponding to actual VM domain] Only done as part of plotting PGA, PGV, PSA, PSA RATIOS Where to get vs30 values?

Obtain observations from GeoNet		A	Took 5 mins to create input file for downloading data from geoNet
(a) Get recorded motion from		h S	Finished downloading data in 13.1 secs
(b) Dresses and refermet using		an	Done Vs30 calcs in 25.0 secs
defaults			Processing .V1A files
(c) Produce plots of acc/vel time			Done processing in 219.4 secs
(d) Droduce IM dete files for			Creating acc and vel plots for all Observed SMSs
stations on map plots in GMT			Done plotting acc, vel in 160.6 secs.
			Creating pSA plots for all Observed SMSs
(e) Produce Vs30 input files for			Done pSA plots in 181.2 secs.
stations			Writting IMs for GMT plotting
			Done in 32.2 secs.
			python2 run_event.py 614.36s user 5.28s system 98% cpu 10:32.05 total
			10:32 to 11:11 to run post processing code with observations only. code ran in
			192.43s, took 5 mins to set the input file. The inputs for this input file
			and the input file for downloading data are duplicated.
			using a log scale for plotting observations only IMs on map results in values smaller than 1 to be
			negative and difficult to interpret. Had to manually adjust cpt color scales for
			better plots, and manually move gmt plots otherwise they are over written.
Plot IM values at GeoNet stations on map (for various IMs)	3 mins	Vi kt or	Viktor was away, and I (Ahsan) performed this task. It was some what unclear how the GMT plotting scripts (behind the scenes) got its input, e.g the domain boundary. The code ran without problems. There was a need to display the color bar on a log scale, which I was not able to do through the GMT scripts but had to rewrite the natural log of the data instead of the plain data. It was unclear what the log=True option achieves in the python wrapper to the GMT scripts.
Plot observed vel time series on	5mins	Vi	skipped, need to finish script that does this
пар		or	
Plot IM values as a function of Rrup and compare with empirical ground motion model (various IMs)	10 mins	A h s an	This was still performed with groundMotionSationsAnalysis code by passing Obs data twice (as Obs and Sim data). This task will be performed along side task (c).
4. Perform GM Simulation (SUNG)		
Submit GM Simulation			(1) We need a standardized event name stored somewhere.
(a) prepare input params.py, pointing to SRF, VM, and stations	(a)5mins	S u	(2) File transfer to Fitzroy could be more simplified. Occasionally I had no right permission to access the files generated by other team members.
of interest etc. (gen_cords.sh and- statlist2gp.py need to be run)- File transfer and install	(b)1:40~3: 30 hours (queue+run)	ng	==> Solution: Each module (VM, Source, Station) can have an "install" script that is executed by the person-in- charge that transfers the necessary files to the appropriate location (derived from the standardised event name)
(b) LF sim (queue time + run time)	(c)13 min		(3) Many codes from various repositories are increasingly becominginter-dependent. We need to make a shared library soon.
(c) Once LF sim complete, run HF and BB sim			(4) Wall clock limit estimation is reasonably accurate, with Fitzroy being prone to fluctuate occasionally. (IO suspected). DB needs some tweak to mark outliers.
5. Post-processing (VIKTOR)			
LF Time slice animation	9 min		1) The animation is working fine for single event. However, if there are 2 SRF files in the run, you need to modify
(a) tar, download data from Fitzroy to local		Vi kt	the parameter file such that it will generate the animation of the specific event that you want2) we might want to generate the animation for the multiple events as well??
(b) From TS produce PGV and MMI plot		or	3) It would be great if we could include "*modelparams*" in the tar file from Fitzroy, which is needed for (5f) plotting
(c) Produce TS animation			

Script preparation and test At SMS locations compare sim vs. obs	10mn (a)–(d) 20 mins	H o by	I have still used Richard code because Daniel/Jason mentioned that they sill have some issue in the std of bias between Obs and empiricalGMM. Once the PP is done there will be significant save in time in terms of file preparation because I will use the same config. as Ahsan
(a) compare waveforms			
(b) Compare IMs as a function of Rrup (multiple IMs; sim, obs, empirial)			
(c) Compare sim, obs, empirical response spectra at individual locations			
(d) Plot overall bias as a function of vibration period (both sim and empirical)			
(e) Preparing input for (f)			
(f) Plot bias(residual) at specific locations on a map	(e) 4 mins (f) 5 mins		(f) requires modelparamsfile/Srf; need to make sure we have them copied in hypocentrer

	Tasks	Time estimation
Hoby	1) Srf plot location need to be modified	2days
	2)	2days
	2)	2-5 days
Ethan	 Streamline the VM generation with gen_coords.csh Write scrip to stitch together multiple PDFs in PDFtk VM domain baseplot to be passed to plot SRF by others 	2days (with other minor tasks)
Ahsan	(1) The input file for post processing and the input file for downloading data are duplicated. Need to fix this duplication on my part.	(1) 2 hours
Sung	 The following 3 items are easy to automated, and where we can easily make mistakes. Create shared parameter files that can be imported into other steps in the workflow (Saves need for configuring params.py and get_cords.sh) Automated file transfer to Fitzroy Automated installation of a job 	1. 2 days 2. 2 days 3. 2 days
Viktor	 Finish script to plot observed timeseries on map Run plots automatically when data has been retrieved Determine format and implement for plotting stations on map with PGV, PGA, PSA data etc and figure out easiest way to plot PGV stations on overlay 	2.5 days 1 - 2 days 2 days
Daniel	 Non-uniform grid should probably run after the VM Include non-uniform grid generation to the automatized workflow Parameters for the grid should be read from the VM file Make sure the format of the output is usable 	

1. Feb 2017 drill (2007 Gisborne Earthquake)

Task	Expected time reqd.	P e o	Notes
4.0		pie	
1. Source (HOBY)			
GeoNet Mw email to SRF			a) Manually extract location, dip,strike, rake from Mw tensor (Ristau) into setSrtParams.py
(a) from Mw tensor get Rup parameters	1minute	H O	b) Change TYPE=2 (point source to finite fault)
(b) From Rup parameters use createSRF.py to make rupture SRF	5mins	by	c) Run "createSRF.py" (Make sure to use python2)
(c) Execution of createSRF.py	10sec		
	6min total		
Plotting of SRF			
(a) SRF plane(s) standard plotting	3mins	н	(a) change manually the SRF and output name in plotSRF.py (Viktor is currently working in improving this part)
(b) SRF on map to verify location (to come after domain set in step 2)	5mins	o by	Execute 'plotSRF.py' (Make sure to use python2)
	2mins		(b) plot_srf.py file.srf
	10min total		
2. Velocity Model Domain (ETHAI	N/ROBIN)		
From rupture location to VM dev and domain	1mins 20 mins	Et h	 Tidy up params_vel.py (some params are redundant) and include corners for plotting before mesh generation
(a) From location and Mw determine VM origin, 3D extent, and simulation duration	(on multiple cores)	n, R o	 Run gen_corrds during VM and place all model params alongside VM files (Done) Integrate the gen_cords into the first step to enable the domain to be plotted on a map and checked / changed if necessary (Done)
(b) Plot VM domain extent	15 mins for	bi n	 Split the VM generation from VM interrogation / slice generation to allow the VM to be copied while slices are being generated
(c) Generate V/M with H=0.1km for	slices		5. Will need a wall clock time estimation done before VM calculation (Done)
this domain, then extract slices	4 mins for		 7. Change names of to be consistent ORIGIN_LAT, ORIGIN_LON,ORIGIN_ROT retain these names , (MODEL_LON_DOT) (Dece)
	VM		8. Write xy2ll into the gen_params.py script to write the domain corners into params_vel.py (Done)
	creation (30 mins for H=0.		 Plot the vm domain box within gen_params.py Include gen_cords.py within genDomain to keep coords and .p .s and .d binary files in together (Done Duplicate of 2)
	20min		There was an error in the auto-VM generation on hypocentre which needed to be fixed, and then the code had to be obtained from github again. (Done)
	totai		(Sung) == items below have beem completed after Feb. drill
			1. nx, ny, nz are automatically updated if user manually edits other values like (extent_x,_y,_zmax)
			2. if extent_x, extent_y and extent_zmax are not divisible by hh, it automatically increases extent_* to the first divisible value
			3. it will automatically work out flo (based on min_vs/(5*hh). The initial value of flo is automated, but user can manually override it by editing params_vel.py (More in item 4)
			4.In params_vel,py, extent_x, extent_y, and extent_zmax, sim_duration and flo have automatically generated initial value. User can manually edit params_vel.py to override them, but user MUST run *validate_parameters. py*, which will check the integrity of the parameter file, and adjust automated values such as nx, ny, nz and sufx. This will overwrite params_vel.py (also keeps a backup of old params_vel.py) and show before and after changes side-by-side.
			(Ethan)
			Remove topography from preliminary plot to speed up investigateDomain.py
			Change default CPTs to extend their range
From non-uniform station grid, get			
statgrid points	(ignore for		
	now)	A	[need .II file for all of NZ (Ahsan)] (Done)
(b) Locations of GeoNet strong motion stations	1-2 hours!!	h s	run statcords (to get station locations corresponding to actual VM domain]
(c) From NZ Vs30 model get Vs30	10mins	a n,	
values at all points	<u>~2hours</u>	C hr	
		is	

(a) Plot VM domain extent, stations (statgrid and GeoNet),(b) color based on Vs30 values	2mins	(a) Vi kt or (b) N ot d o ne	Only done as part of plotting PGA, PGV, PSA, PSA RATIOS Where to get vs30 values?
3. Observed strong motions (SUN	G)		
Obtain observations from GeoNet (a) Get recorded motion from GeoNet ftp (b) Process and reformat using defaults (c) Produce plots of acc/vel time series, response spectra (d) Produce IM data files for stations on map plots in GMT (e) Produce Vs30 input files for stations	(a) to (d) took 7.5 mins. (e) took 40 mins	A h s an	 Tasks (a) to (d) are automated but (e) is still manual and took the longest time. In (e) the computational time itself only took about 2 mins, it was the formatting that took the most time. 2017-02-23_133554 Mw4.7 simulated as a point source. Since the earthquake occurred within the FVM domain, no new velocity model was generated. The three key steps (in absence of observed data) were (1) generating the point source (srf file) (2) Running the simulation (3) Producing time slice maps/movie, PGV and MMI maps. The main takeaway lessons were that (1) Generating a velocity model may not be necessary for each simulation drill (2) A need for quickly deciding whether a new velocity model should be created. Show existing domain on Map!? (3) May need to generate file names and directory structure at the time of source generation for efficiency.
Plot IM values at GeoNet stations on map (for various IMs)	3 mins	Vi kt or	Viktor was away, and I (Ahsan) performed this task. It was some what unclear how the GMT plotting scripts (behind the scenes) got its input, e.g the domain boundary. The code ran without problems. There was a need to display the color bar on a log scale, which I was not able to do through the GMT scripts but had to rewrite the natural log of the data instead of the plain data. It was unclear what the log=True option achieves in the python wrapper to the GMT scripts.
Plot observed vel time series on map	5mins	Vi kt or	skipped, need to finish script that does this
Plot IM values as a function of Rrup and compare with empirical ground motion model (various IMs)	10 mins	A h s an	This was still performed with groundMotionSationsAnalysis code by passing Obs data twice (as Obs and Sim data). This task will be performed along side task (c).
4. Perform GM Simulation (SUNG)		
Submit GM Simulation (a) <u>prepare input params.py,</u> peinting to SRF, VM, and stations of intercest etc. (gon_eords.sh and- statlist2gp.py need to be run). File transfer and install (b) LF sim (queue time + run time) (c) Once LF sim complete, run HF and BB sim	(a)5mins (b)1:40~3: 30 hours (queue+run) (c)13 min	S u ng	 (1) We need a standardized event name stored somewhere. (2) File transfer to Fitzroy could be more simplified. Occasionally I had no right permission to access the files generated by other team members. =>> Solution: Each module (VM, Source, Station) can have an "install" script that is executed by the person-incharge that transfers the necessary files to the appropriate location (derived from the standardised event name) (3) Many codes from various repositories are increasingly becoming inter-dependent. We need to make a shared library soon. (4) Wall clock limit estimation is reasonably accurate, with Fitzroy being prone to fluctuate occasionally. (IO suspected). DB needs some tweak to mark outliers.
5. Post-processing (VIKTOR)			
LF Time slice animation (a) tar, download data from Fitzroy to local (b) From TS produce PGV and MMI plot (c) Produce TS animation	9 min	Vi kt or	 The animation is working fine for single event. However, if there are 2 SRF files in the run, you need to modify the parameter file such that it will generate the animation of the specific event that you want we might want to generate the animation for the multiple events as well?? It would be great if we could include "*modelparams*" in the tar file from Fitzroy, which is needed for (5f) plotting

Script preparation and test	10mn	H o	It would be great to have some systematic way to select the "good" waveforms (Probably in the Observation data processing??) and have new station. If corresponding to these acceptable Obs waveform for validation
At SMS locations compare sim vs. obs	(a)–(d) 20 mins	by	F
(a) compare waveforms			
(b) Compare IMs as a function of Rrup (multiple IMs; sim, obs, empirial)			
(c) Compare sim, obs, empirical response spectra at individual locations			
(d) Plot overall bias as a function of vibration period (both sim and empirical)			
(e) Preparing input for (f)			
(f) Plot bias(residual) at specific	(e) 4 mins		
ισσαιιστις στι α πιαρ	(f) 5 mins		(f) requires model params file/Srf; pand to make sure we have them capied in hyperaptran

	Tasks	Time estimation
Hoby	1) Include ground motion duration in the PP plotting	2days
	2) Add the empirical ground motion duration	2days
	2) get plotSRF improved	2-5 days
Ethan	 Streamline the VM generation with gen_coords.csh Write scrip to stitch together multiple PDFs in PDFtk VM domain baseplot to be passed to plot SRF by others 	2days (with other minor tasks)
Ahsan	(1) automate process to produce Vs30 files and their formatting.	(1) 2 hours
	(2) Produce plots that need Rrup within the existing geoNet real-time processing scripts as opposed to separately	(2) 2 hours
	round Motion Stations Analysis code separately	(3) 2 hours
	(3) Include header information for IM data files to be plotted on map using GMT. Currently this is done manually.	(4) 2hours
	(4) Incorporate change to write custom header to .000 .090 .ver GP format files. Issue brought up by Chris McGann	
Sung	 The following 3 items are easy to automated, and where we can easily make mistakes. 1. Create shared parameter files that can be imported into other steps in the workflow (Saves need for configuring params.py and get_cords.sh) 2. Automated file transfer to Fitzroy 3. Automated installation of a job 	1. 2 days 2. 2 days 3. 2 days
Viktor	 Finish script to plot observed timeseries on map Run plots automatically when data has been retrieved Determine format and implement for plotting stations on map with PGV, PGA, PSA data etc and figure out easiest way to plot PGV stations on overlay 	2.5 days 1 - 2 days 2 days
Daniel	 Non-uniform grid should probably run after the VM Include non-uniform grid generation to the automatized workflow Parameters for the grid should be read from the VM file Make sure the format of the output is usable 	

1. Jan 2017 drill (Lake Grassmere Earthquake)

Task	Expected time	P e	Notes
	reqd.	o ple	
1. Source (HOBY)			
GeoNet Mw email to SRF			a) Manually extract location, dip,strike, rake from Mw tensor (Ristau) into setSrfParams.py
(a) from Mw tensor get Rup	1minute	Н	b) Change TYPE=2 (point source to finite fault)
(b) From Rup parameters	5mins	by	c) Run "createSRF.py"
use createSRF.py to make rupture SRF			
(c) Execution of createSRF.	10sec		
pý	6min total		
Plotting of SRF			
(a) SRF plane(s) standard	3mins	Н	(a) change manually the SRF and output name in plotSRF.py (need to be improved)
(b) SRF on map to verify	5mins	by	Execute 'plotSRF.py'
location (to come after domain set in step 2)	2mins		(b) plot_srf.py file.srf
,	10min total		
2. Velocity Model Domain (E	THAN/ROB	IN)	
From rupture location to VM	1mins	Et	1. Tidy up parame, yel by (some parame are redundant) and include corners for plotting before mesh generation
(a) From location and Mw	20 mins	a	 Run gen_corrds during VM and place all model params alongside VM files Integrate the gen_corrds into the first step to enable the domain to be plotted on a map and checked / changed if
determine VM origin, 3D extent, and simulation	multiple cores)	R	necessary 4. Split the VM generation from VM interrogation / slice generation to allow the VM to be copied while slices are
duration	, 15 mins	bi n,	being generated 5. Will need a wall clock time estimation done before VM calculation
(b) Plot VM domain extent	for slices		 Merge EXTENT_Z_SPACING, EXTENT_LATLON_SPACING and HH Change names of to be consistent ORIGIN_LAT, ORIGIN_LON, ORIGIN_ROT retain these names ,
(c) Generate VM with H=0.1km for this domain, then	4 mins for H=0.		(MODEL_LAT,_LON,_ROT) 8. Write xy2ll into the gen_params.py script to write the domain corners into params_vel.py
extract slices	2km VM creation		 Plot the vm domain box within gen_params.py Include gen_cords.py within genDomain to keep coords and .p .s and .d binary files in together
	(30 mins for H=0.		
	1km)		1. There was an error in the auto-VM generation on hypocentre which needed to be fixed, and then the code had to
	20min total		be obtained from github again.
From non-uniform station grid, get statgrid points			
(a) Locations of statgrid	(ignore for now)		
points	1-2	A h	[need .ll file for all of NZ (Ahsan)] (Done)
(b) Locations of GeoNet strong motion stations	hours!!	s a	run statcords (to get station locations corresponding to actual VM domain]
(c) From NZ Vs30 model get	10mins	n, C	
Vs30 values at all points	<u>~2hours</u>	hr is	
(a) Plot VM domain extent, stations (statorid and	2mins	(a	Only done as part of plotting PGA, PGV, PSA, PSA RATIOS
GeoNet),		/ Vi kt	Where to get vs30 values?
(b) color based on Vs30 values		or	
		(b)	
		N ot	
		d o	
		ne	

3. Observed strong motions	3. Observed strong motions (SUNG)			
Obtain observations from GeoNet (a) Get recorded motion from GeoNet ftp (b) Process and reformat using defaults (c) Produce plots of acc/vel time series, response spectra	45 mins total for (a) and (b) 60 mins for generatin g Vs30 values. (one-off task) 5 mins for (c)	A h s an	Majority of time spent trouble shooting. See comment below GeoNet people make errors which can cause errors and breakdown our code. The data location has files that should have been removed. Take a look at ftp://ftp.geonet.org.nz/strong/processed/Proc/2013/08_Aug/2013-08-16_023105 /Vol1/data . Files ending with _ should not be there and nor should V2A and V3A data be there. You will also see three stations named like Q13924J03.V1A that don't follow the naming convention used for all the other data files. This broke down my code. For the next drill, I will change the code to catch these as exceptions and continue smoothly.	
Plot IM values at GeoNet stations on map (for various IMs)	3 mins	A h s an	Calculations of PGVs, PGAs and pSA for set of chosen periods took about 3 mins, these were then passed on to Viktor who plotted them.	
Plot observed vel time series on map	5mins	Vi kt or	skipped, need to finish script that does this	
Plot IM values as a function of Rrup and compare with empirical ground motion model (various IMs)	10 mins	A h s an	Code ran as expected.	
4. Perform GM Simulation (S	UNG)			
Submit GM Simulation (a) <u>prepare input parame.py</u> , <u>pointing to SRF, VM, and</u> <u>stations of interest otc.</u> (gen_cords.sh and statlist2gp. <u>py need to be run)</u> File transfer and install (b) LF sim (queue time + run time) (c) Once LF sim complete, run HF and BB sim	(a)5mins (b)5 hours (queue+r un) (c)13 min	S u ng	gen_ts was broken (fixed) Will be nice to have auto file transfer (in 3 separate streams esp. for VM model) Will need to separate HF from BB and run it alongside LF.	
5. Post-processing (VIKTOR)				
LF Time slice animation (a) tar, download data from Fitzroy to local (b) From TS produce PGV and MMI plot (c) Produce TS animation	10mins 25mins 25min	Vi kt or	needed to fix up scripts to work with new parameter style mainly needed to fix script to use new parameter style, should run automatically after step A run time only ~3 minutes need to automate a little more, run time ~5 mins	

Script preparation and test	10mn	H	
At SMS locations compare sim vs. obs	(a)–(d) 30 mins	by	 Testing the Script using only the observation and remove irrelevant stations Make sure to use fd_*.ll station lists instead of geonet*.ll Need to add Ground motion duration
(a) compare waveforms			
(b) Compare IMs as a function of Rrup (multiple IMs; sim, obs, empirial)			
(c) Compare sim, obs, empirical response spectra at individual locations			
(d) Plot overall bias as a function of vibration period (both sim and empirical)			default parameters used, runs in 2 mins but needed to fix a few things up
(e) Preparing input for (f)			
(f) Plot bias(residual) at specific locations on a map	(e) 4 mins		
	(f) 5 mins		
		(f)	
		ΓL	

	Tasks	Time estim ation
Hoby	1) Include ground motion duration in the PP plotting	2days
	2) Add the empirical ground motion duration	2days
	2) get plotSRF improved	2-5 days
Ethan	 Streamline the VM generation with gen_coords.csh Write scrip to stitch together multiple PDFs in PDFtk VM domain baseplot to be passed to plot SRF by others 	2days (with other minor tasks)
Ahsan	 (1) GeoNet people make errors which can cause errors and breakdown our code. The data location has files that should have been removed. Take a look at ftp://ftp.geonet.org.nz/strong/processed/Proc/2013/08_Aug/2013-08-16_023105/Vol1/data . Files ending with _ should not be there and nor should V2A and V3A data be there. You will also see three stations named like Q13924J03.V1A that don't follow the naming convention used for all the other data files. This broke down my code. For the next drill, I will change the code to catch these as exceptions and continue smoothly. (2) Deal with new stations better. Currently I maintain a list of about 1350 stations and new stations coordinates are read from there. But in this drill one new station was not in that extensive list. A better solution would be to read the data file and extract the coordinates rather than relying on a list. (3) I shall add Vs30 = 500 for any new stations and give the list of new stations to Kevin after our drills. He can then provide a better Vs30 estimate. This way we simply the problem of knowing Vs30 for new stations. What do you propose? These are not big issues in general but they are if we want unbroken, smooth work flow during real time simulation. I shall add these to my task list. 	(1) 2 hours (2) 5 hours (3) 2 hours

Sung	 The following 3 items are easy to automated, and where we can easily make mistakes. 1. Create shared parameter files that can be imported into other steps in the workflow (Saves need for configuring params.py and get_cords.sh) 2. Automated file transfer to Fitzroy 3. Automated installation of a job 	1. 2 d ys 2. 2 d a ys 3. 2 d a ys 3. 2 s
Viktor	 Finish script to plot observed timeseries on map Run plots automatically when data has been retrieved Determine format and implement for plotting stations on map with PGV, PGA, PSA data etc and figure out easiest way to plot PGV stations on overlay 	2.5 days 1 - 2 days 2 days
Daniel	 Non-uniform grid should probably run after the VM Include non-uniform grid generation to the automatized workflow Parameters for the grid should be read from the VM file Make sure the format of the output is usable 	

1. Dec 2016 drill (Wilberforce Earthquake)

Task	Expected time reqd.	People	Notes
1. Source (HOBY)			
GeoNet Mw email to SRF (a) from Mw tensor get Rup parameters (b) From Rup parameters use createSRF.py to make rupture SRF (c) Execution of createSRF.py	1minute 5mins 10sec 6min total	Hoby	 a) Manually extract location, dip,strike, rake from Mw tensor (Ristau) into setSrfParams.py b) Change TYPE=2 (point source to finite fault) c) Run "createSRF.py"
Plotting of SRF			Execute 'plotSRF.py' (select title for top of script)
(a) SRF plane(s) standard plotting	3mins	(a)	(a) Hoby : Just needed to change SRF name
(b) SRF on map to verify location (to come after domain set in	5mins	(b) Viktor	(b) Viktor: make 2 plots on 1 image, 1 zoomed in with slip visible, 2 whole
siep 2)	2mins		
	10min total		Finished automating. only SRF needed, no reliance on params.py etc
2. Velocity Model Domain (ETHAN/ROBIN)			·
From rupture location to VM dev and domain		Ethan,	1) Origin lat/lon/depth written into 'generate_parameters.py' python script
(a) From location and Mw determine VM origin, 3D extent, and simulation duration	2mins		2) Run 'generate_parameters.py', output is a text file which is read by the next step
(b) Plot VM domain extent	1minc		3) Run 'genDomain.py' this generates the velocity model, extracts slices.
(c) Generate VM with H=0.1km for this domain			
15 mins multiple	multiple cores)		Need to implement duration calculations
	20min total		Need to streamline Gen_Coords to sync with model parameters
			Run gen_cords.csh and plot
			Create .p/s/d files

From non-uniform station grid, get statgrid points				
(a) Locations of statgrid points	(ignore for now)			
(b) Locations of GeoNet strong motion stations	1-2 hours!!	Ahsan	[need .ll file for all of NZ (Ahsan)] (Done)	
(c) From NZ Vs30 model get Vs30 values at all points	10mins	, Chris	run statcords (to get station locations corresponding to actual VM domain]	
	~2hours			
(a) Plot VM domain extent, stations (statgrid and GeoNet),	2mins	(a)		
(b) color based on Vs30 values		VIKTOR	Where to get vs30 values?	
		Not		
2. Observed streng metians. (SUNC)		done		
3. Observed strong motions (SUNG)	40 mine tetal	Ahaan		
(a) Optimized matter from Cookiet	40 mins totai	Ansan	Uploading to cloudup was tricky as the files need to be downloaded from	
(a) Get recorded motion from Georee up			the linux dox first.	
(b) Process and reformat using defaults				
(c) Produce plots of acc/vel time series, response spectra				
Plot IM values at GeoNet stations on map (for various IMs)	1 hour (GMT)	Ansan	updating e3d.par domain into	
Plot observed vel time series on map	5mins	Viktor	need to automate location of stations involved, zoom levels etc code exists, needs to be brought together / cleaned up	
Plot IM values as a function of Rrup and compare with	20min	Ahsan	Richard's code runs for 3 mins. Had to go through a number of iterations to get nice looking plots	
	1hour 30mins		Has a good idea to reduce the time significantly. Knows what parameters	
			to change.	
			Not easy to automate.	
4. Perform GM Simulation (SUNG)		-		
Submit GM Simulation				
(a) prepare input params.py, pointing to SRF, VM, and	20mins	Sung	1. Hoby uploaded SRF to Fitzroy	
to be run)	512 cores		 Sung opposited a racine for the drill with persons by heving all the 	
(b) LF sim (queue time + run time)	(onode) [wait+run		 Sung cleated a recipe for the drift with parameters by having all the parameters set, and other files copied from 2010Sep sim. 	
(c) Once LF sim complete, run HF and BB sim			4. Modification to install.sh was not necessary, but we will eventually need to.	
	30mins		 Almost forgot to produce .startcords from .ll. For HF, it is advised to use the FD version of .ll (ie. output of statlist2gp.py) in params.py 	
	4nours		Good opportunity for automation. (a) can be done in a sec.	
5. Post-processing (VIKTOR)		1	·	
LF Time slice animation				
(a) tar, download data from Fitzroy to local	30mins	Viktor	(can be streamlined, at later date)	
(a) Produce TS animation	30mins		vm domain used by default, determine if regional sites should be	
(b) From TS produce PGV plot	5min		added (crowding), padding around vm based on rot, domain if rotation small	
(c) Produce MMI plot from PGV 5min			mostly automated TS animation, some parameters needed such as	
			single script does both PGV, MMI - need to finish automating colour range for PGV	

Script preparation	10mn	Hoby	
At SMS locations compare sim vs. obs	(a)–(c) 30 mins		1. Make sure to use fd_*.Il station lists instead of geonet*.Il
(a) compare waveforms			that part of the code to plot the IMs
(b) Compare IMs as a function of Rrup (multiple IMs; sim, obs, empirial)	(d) 20mp		
(c) Compare sim, obs, empirical response spectra at individual locations	(u) 201111		
(d) Plot overall bias as a function of vibration period (both sim and empirical)	(a) 40 mins		
(e) Preparing input for (f)			make a script that plots just this and loops over period values - done self
(f) Plot bias(residual) at specific locations on a map			contained input me used and model_params

	Tasks	Time estimation
Hoby	Fix bias plotting for both empirical and physics-based models	2days
Ethan	 Streamline the VM generation with gen_coords.csh Write scrip to stitch together multiple PDFs in PDFtk VM domain baseplot to be passed to plot SRF by others 	2days (with other minor tasks)
Ahsan	 Update all_geoNet_stations.II to include SMS that closed in 2010. This will add more stations to the list and we won't have a problem when dealing with historical events. Kevin would need to update the Vs30 stations list accordingly. Some python scripts (code that downloads data, code that processes data) that run one after another can and will be automated. Coordinates with Viktor for plotting IMs on map. Currently I and Viktor separately gather velocity model domain data etc, for GMT plotting, which is a manual task. Supply Viktor IMs data file for him to plot. 	5days
Sung	 The following 3 items are easy to automated, and where we can easily make mistakes. 1. Create shared parameter files that can be imported into other steps in the workflow (Saves need for configuring params.py and get_cords.sh) 2. Automated file transfer to Fitzroy 3. Automated installation of a job 	 1. 2 days 2. 2 days 3. 2 days
Viktor	 Fix plotting script for SRF to show fault location clearly, Create plotting script that takes PGV and MMI, create plotting script for station resuduals. Automatically calculate default parameters where possible, allow scripts to be run where paths / needed inputs can be passed to them. connect scripts where input from one should automatically run another where possible eg: create PGV, MMI data and instantly plot it 	1 - 2 days 1 - 2 days 1 - 2 days
Daniel		